

# PRODUCT SPECIFICATION



## HYB 131-10 Anti-MBL (human)

### *Mouse monoclonal antibody*

OVERVIEW	Article No.	100884 (0.2 mL), 100885 (1.0 mL)	
	Product Name	HYB 131-10 Anti-MBL (human)	
	Clone ID	1E2	
	Subclass	IgG1 / Kappa	
	Specificity	HYB 131-10 Anti-MBL (human) is specific for MBL from human serum or plasma.	
	Species Reactivity	Human	
	Epitope Specificity	The epitope specificity differs from that of HYB 131-01 and HYB 131-11.	
	Immunogen	MBL purified from human donor plasma.	
	Fusion Partner	X63-Ag8.653.	
	Culture Medium	Dulbecco's modified Eagle's medium with 10 % fetal calf serum	
TESTED APPLICATION	Method	Usability	References
	Enzyme linked immunosorbent assay (ELISA)	Yes	In house analysis, 1
	Western Blot (WB)	Yes	In house analysis
PRODUCT SPECIFIC INFORMATION	<p>In ELISA HYB 131-10 reacts strongly with MBL coated directly onto the well. HYB 131-10 reacts with MBL in its oligomerized form and as a single subunit of 26 kDa. HYB 131-10 can be used in sandwich ELISA in combination with HYB 131-11 for quantitation of total MBL.</p> <p>In Western blotting, HYB 131-10 reacts with human MBL both in its polymeric conformation and as a single subunit of app. 26 kDa.</p>		
PROPERTIES	Conjugation:	Unconjugated	
	Form	Liquid	
	Preparation:	Protein A	
	Concentration:	1 mg/mL $\pm$ 10%, based on A <sub>280</sub> . See Certificate of Analysis for details.	
	Solvent:	PBS, pH 7.2 – 7.4	
	Storage information:	Store at $\leq$ - 18 °C.	

### SSI Antibodies

Statens Serum Institut, 5 Artillerivej, DK-2300 Copenhagen S, Denmark

E [ssi-antibodies@ssi.dk](mailto:ssi-antibodies@ssi.dk) | W <https://antibodies.ssi.dk/> | CVR/VAT DK-46837428

# PRODUCT SPECIFICATION



<b>TARGET</b>	<p>Mannan-binding lectin (MBL), also called mannose-binding lectin or protein, is a C-type lectin and an important component in innate immunity. MBL is an oligomer i.e. forming dimers to hexamers of homotrimeric subunits of approximately 26 kDa polypeptides. This oligomerization is essential for functional activity (2).</p> <p>MBL forms a non-covalent complex with specific MBL-associated serine proteases (MASPs), termed MASP-1, -2, and -3. Upon binding to the surface of a pathogen, MASP-activation is initiated with subsequent complement activation and clearance through lysis or phagocytosis (3).</p> <p>MBL-deficiency is the most common immune defect resulting in susceptibility to severe infections in early childhood, or if immuno-suppressed (4). MBL-deficiency has also been associated with several clinical disorders, e.g. autoimmune diseases, endocarditis, and septicaemia (4,5).</p> <p>Normal levels of oligomeric MBL in serum are 1 – 5 µg/mL whereas MBL-deficient serum levels are &lt; 100 ng/mL, when estimated by a standard ELISA for MBL quantification (3). Due to the presence of different structural and promotor alleles 12 % or more of the Caucasian population have low concentrations (&lt; 50 ng/mL) of normally oligomerized, functional MBL in plasma or serum (6).</p>
<b>REFERENCES</b>	<ol style="list-style-type: none"><li>1) US patent No. US 7,211,396 B2</li><li>2) Laursen I, Hojrup P, Houen G, Christiansen M. (2008) Characterization of the 1st SSI purified MBL standard. Clin Chim Acta, 395(1-2), 159-161.</li><li>3) Dommett RM, Klein N, Turner MW. (2006) Mannose-binding lectin in innate immunity: past, present and future. Tissue Antigens, 68(3):193-209.</li><li>4) Kilpatrick DC. (2002) Mannan-binding lectin: clinical significance and applications. Biochim Biophys Acta, 1572(2-3): 401-413.</li><li>5) Tran CT, Kjeldsen K, Haunsø S, Høiby N et al. (2007) Mannan-binding lectin is a determinant of survival in infective endocarditis. Clin Exp Immunol, 148(1): 101-105.</li><li>6) Steffensen R, Thiel S, Varming K, Jersild C, Jensenius JC (2000) Detection of structural gene mutations and promoter polymorphisms in the mannan-binding lectin (MBL) gene by polymerase chain reaction with sequence-specific primers. J Immunol Methods 241:33-42.</li></ol>

Version 1 · April 2022

## Conditions

### **For research use only. Not for use in diagnostic procedures. Not for therapeutic use or applications.**

*The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall Statens Serum Institut be responsible for loss of profits or indirect consequential losses resulting from use of its products. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.*

## **SSI Antibodies**

Statens Serum Institut, 5 Artillerivej, DK-2300 Copenhagen S, Denmark

E [ssi-antibodies@ssi.dk](mailto:ssi-antibodies@ssi.dk) | W <https://antibodies.ssi.dk/> | CVR/VAT DK-46837428