

PRODUCT SPECIFICATION

STATENS
SERUM
INSTITUT



HYB 123-16 Anti Diphtheria toxin (DT)

Mouse monoclonal antibody

Article No.	66046 (0.2 mL), 101052 (1.0 mL)	
Product Name	HYB 123-16 Anti Diphtheria toxin (DT)	
Clone	6B5	
Subclass	IgG1 / Kappa	
Description	<p>Preparation: Protein-A purified</p> <p>Concentration: 1 mg/mL \pm 10%, based on A₂₈₀. See Certificate of Analysis for details.</p> <p>Solvent: PBS, pH 7.2 – 7.4</p> <p>Storage: -18 °C or colder</p>	
Antigen	<p>Diphtheria toxin (DT) is secreted by certain strains of <i>Corynebacterium diphtheriae</i> and catalyzes the ADP-ribosylation of eukaryotic aminoacyl-transferase II (EF-2) using NAD as a substrate [1]. This reaction forms the basis for its toxicity towards eukaryotic organisms [2].</p> <p>Diphtheria toxin is synthesized and excreted as a proenzyme, composed of a single polypeptide chain having a molecular weight of approximately 63 kDa [3]. Two covalent alterations in structure are necessary for expression of its enzymatic activity. First, mild proteolysis results in the formation of "nicked toxin", which is enzymatically inactive and consists of two major fragments, A and B, linked by a disulfide bond. Reduction of the nicked toxin with thiols releases the N-terminal A fragment (~24 kDa) which is enzymatically active.</p> <p>The C-terminal B fragment (~39 kDa) has no apparent enzymatic activity, but is required for toxicity. Evidence suggests that the B fragment is responsible for recognizing and binding the toxin to cell surface receptors [4].</p>	
Immunogen	Diphtheria toxoid (formaldehyde inactivated Diphtheria toxin).	
Specificity	HYB 123-16 reacts primarily with Diphtheria toxin with a low cross reactivity with Diphtheria toxoid.	
Epitope Specificity	HYB 123-16 reacts with a different epitope compared to HYB 123-09 and HYB 123-18.	
Reactivity	HYB 123-16 reacts well in ELISA with coated Diphtheria toxin. When used in WB HYB 123-16 detects a band at approximately 63 kDa corresponding to the proenzyme.	
Culture Medium	Dulbecco's modified Eagle's medium with 10 % FCS.	
Fusion Partner	X63-Ag8.653.	
Immunization	Female CF1xBalb/c F1 hybrid mice were immunized i.p. with immunogen	
Application	Method	Usability
	ELISA	yes
	Immunoblotting	yes
	Immuno.fluoresc.	nd.

See next page

HYB 123-16 Anti Diphtheria toxin (DT)

mouse monoclonal antibody

STATENS
SERUM
INSTITUT



References

- 1) **Honjo, J., Nishizuka, Y., Hayaishi, O. and Kato, I.** (1968) J. Biol. Chem. 243, 3553-3555.
- 2) **Gill, D.M., Pappenheimer, Jr., A.M. and Baseman, J.B.** (1969) Cold Spring Harbor Symp. Quant. Biol. 34, 595-602.
- 3) **Collier, R.J. and Kandel, J.** (1971) J. Biol. Chem. 246, 1496-1503.
- 4) **Ittelson, T.R. and Gill, D.M.** (1973) Nature 242, 330-332.

Conditions

For research use only. Not for use in diagnostic procedures. Not for therapeutic use or applications.

The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall Statens Serum Institut be responsible for loss of profits or indirect consequential losses resulting from use of its products. The animals from which this product was derived have not been exposed to or inoculated with any livestock or poultry disease agents exotic to the United States or Western Europe, and did not originate from facilities where work with exotic disease agents affecting livestock or avian species is carried out.